arXiv:1207.6800v1 [cond-mat.supr-con] 29 Jul 2012

Additional *is*-wave Cooper pairing driven by nematic order in *d*-wave superconductors

Jing-Rong Wang and Guo-Zhu Liu
Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230026, P. R. China

We make a non-perturbative analysis of the strong interaction between gapless nodal fermions and nematic order parameter in two-dimensional $d_{x^2-y^2}$ superconductors. We predict that the critical nematic fluctuation can generate a dynamical nodal gap if the fermion flavor N is smaller than a threshold N_c. Such gap generation leads to an additional *is*-wave Cooper pairing instability, and induces a fully gapped $d_{x^2-y^2}+is$ superconducting dome in the vicinity of nematic quantum critical point. The opening of dynamical gap has important consequences, including saturation of fermion velocity renormalization, weak confinement of fermions, and suppression of observable quantities.

PACS numbers: 73.43.Nq, 74.20.Rp, 74.25.Dw

One of the most prominent properties of high-T_c copper-oxide superconductors is that they exhibit a number of long-range orders upon changing the chemical doping, such as antiferromagnetism, superconductivity, stripe, nematic state, and so on. The competition and coexistence between superconductivity and other long-range orders are believed to be fundamental issues in the studies of high-T_c superconductors. Among the orders that are in competition with superconductivity, the nematic order, which spontaneously breaks C_4 symmetry down to C_2 symmetry, has attracted special theoretical and experimental interest in the past decade [1–4].

In recent years, strong anisotropy in electronic properties has been observed in various experiments performed on $\mathrm{YBa_2Cu_3O_{6+\delta}}$ [5–7] and $\mathrm{Bi_2Sr_2CaCu_2O_{8+\delta}}$ [8]. Such anisotropy is widely attributed to the formation of an electronic nematic state. An important indication of these experiments is the existence of nematic quantum phase transition in these two typical superconductors. Interestingly, similar nematic states are also observed in several other correlated electron systems, including iron-based superconductor [9], heavy fermion superconductor [10], $\mathrm{Sr_3Ru_2O_7}$ superconductor [11], and semiconductor heterostructure [12].

High-T_c superconductors are known to have a $d_{x^2-y^2}$ energy gap, which vanishes linearly at four nodal points $(\pm\frac{\pi}{2},\pm\frac{\pi}{2})$. Due to this property, gapless nodal quasiparticles (qps) are present in the superconducting dome even in the low-energy regime. When a nematic quantum phase transition occurs in the superconducting dome, the fluctuation of nematic order parameter couples to the gapless nodal qps. This coupling becomes singular at the nematic quantum critical point (QCP), and can lead to unusual behaviors. Vojta *et al.* analyzed this model by means of $\epsilon = 3 - d$ expansion and found runaway behavior [13]. Later, perturbative expansion in terms of $1/N$ with N being the fermion flavor has been extensively applied to address this issue [14–20]. For instance, Kim *et. al.* revealed a second-order nematic phase transition after performing a large-N analysis [14]. More recently, renormalization group calculations of Huh and Sachdev [15] discovered a novel fixed point with extreme fermion velocity anisotropy, i.e., $\delta = v_\Delta/v_F \to 0$, where $v_{F,\Delta}$ are Fermi/gap velocities of nodal qps respectively. Subsequent studies showed that such extreme anisotropy can give rise to a variety of nontrivial properties, such as non-Fermi liquid behaviors [16], enhancement of thermal conductivity [17], suppression of superconductivity [19], and first order instability [20].

We should note that all pervious field-theoretic analysis are based on perturbative expansions [13–20]. In this formalism, one starts from the free fermion propagator, $G_0^{-1}(\omega,\mathbf{k}) = -i\omega + v_F k_x\tau^z + v_\Delta k_y\tau^x$, and then calculates the one-loop fermion self-energy $\Sigma(\omega,\mathbf{k})$ that is induced by the interaction with nematic fluctuation. Generically, $\Sigma(\omega,\mathbf{k})$ can be expanded as

$$\Sigma(\omega,\mathbf{k}) = -i\Sigma_0\omega + \Sigma_1 v_F k_x\tau^z + \Sigma_2 v_\Delta k_y\tau^x, \qquad (1)$$

where the functions $\Sigma_{0,1,2}$ are the temporal and spatial components respectively. The fermion damping effect is encoded in Σ_0 [14], whereas the velocity renormalization can be obtained from $\Sigma_{0,1,2}$ [15]. However, in principle there could be the fourth term, $m\tau^y$, corresponding to a nonzero mass gap term of nodal qps. This mass term can never be obtained to any finite order of perturbative expansion, but may be dynamically generated if we perform non-perturbative calculations.

In this paper, we study dynamical gap generation of originally gapless nodal qps due to nematic fluctuation by means of non-perturbative expansion. With the help of Dyson-Schwinger (DS) equation that connects the free and complete propagators of nodal qps, we obtain a nonlinear gap equation of fermion mass m in the vicinity of nematic QCP. After solving this equation, we find that a nonzero mass gap, $m\tau^y$, is dynamically generated when the fermion flavor N is below certain critical value N_c, i.e. $N < N_c$. We demonstrate that the dynamical gap m induced by nematic order corresponds to a secondary *is*-wave Cooper pairing formation, so the critical nematic fluctuation drives a transition from a pure $d_{x^2-y^2}$ superconducting state to a $d_{x^2-y^2}+is$ superconducting state in the vicinity of nematic critical point. As a consequence, the superconducting state is fully gapped and the massive nodal qps are weakly confined by a logarithmic potential. Moreover, the dynamical gap leads to saturation of fermion velocity renormalization and strong suppression of some observable quantities.

We start from the following action [14–19]

$$S = S_\psi + S_\phi + S_{\psi\phi}, \tag{2}$$

where the free action for nodal qps is

$$S_\psi = \int \frac{d^2\mathbf{k}}{(2\pi)^2}\frac{d\omega}{2\pi}\Psi_{1a}^\dagger(-i\omega + v_F k_x\tau^z + v_\Delta k_y\tau^x)\Psi_{1a} + \int \frac{d^2\mathbf{k}}{(2\pi)^2}\frac{d\omega}{2\pi}\Psi_{2a}^\dagger(-i\omega + v_F k_y\tau^z + v_\Delta k_x\tau^x)\Psi_{2a} \tag{3}$$

Here, the Nambu spinor Ψ is defined as, $\Psi_{1a} = (f_{1a}, \epsilon_{ab}f_{3b}^\dagger)^T$ and $\Psi_{2a} = (f_{2a}, \epsilon_{ab}f_{4b}^\dagger)^T$, where $\epsilon_{ab} = -\epsilon_{ba}$ with spin indices a, b. The operators f_1, f_2, f_3 and f_4 are nodal qps excited from four nodal points $(\frac{\pi}{2}, \frac{\pi}{2})$, $(-\frac{\pi}{2}, \frac{\pi}{2})$, $(-\frac{\pi}{2}, -\frac{\pi}{2})$, and $(\frac{\pi}{2}, -\frac{\pi}{2})$, respectively. The free action for nematic order parameter ϕ is

$$S_\phi = \int d^2\mathbf{x}d\tau\left[\frac{1}{2}(\partial\tau\phi)^2 + \frac{c^2}{2}(\nabla\phi)^2 + \frac{r}{2}\phi^2 + \frac{u}{4!}\phi^4\right], \tag{4}$$

where ϕ is a real scalar field since the nematic transition is accompanied by a discrete symmetry breaking (i.e., Ising-type). The coupling between the nodal qps and the nematic order parameter is [13]

$$S_{\psi\phi} = \int d^2\mathbf{x}d\tau\{\lambda\phi(\Psi_{1a}^\dagger\tau^x\Psi_{1a} + \Psi_{2a}^\dagger\tau^x\Psi_{2a})\}. \tag{5}$$

The physical flavor of nodal qps is determined by spin degeneracy, so $N = 2$. However, in order to carry out analytical calculations, it is convenient to assume a large N and then perform $1/N$ expansion. The free propagators for nodal qps Ψ_{1a} and Ψ_{2a} are $G_{\Psi_{1a}}^0(\omega, \mathbf{k}) = (-i\omega + v_F k_x\tau^z + v_\Delta k_y\tau^x)^{-1}$ and $G_{\Psi_{2a}}^0(\omega, \mathbf{k}) = (-i\omega + v_F k_y\tau^z + v_\Delta k_x\tau^x)^{-1}$, respectively. To the leading order of $1/N$ expansion, the polarization function due to particle-hole excitations is given by

$$\Pi(\Omega, \mathbf{q}) = \lambda^2\sum_{a=1}^{N}\sum_{i=1,2}\int\frac{d^2\mathbf{k}}{(2\pi)^2}\frac{d\omega}{2\pi}\mathrm{Tr}\left[\tau^x G_{\Psi_{ia}}^0(\omega, \mathbf{k}) \times\tau^x G_{\Psi_{ia}}^0(\omega + \Omega, \mathbf{k} + \mathbf{q})\right]. \tag{6}$$

After straightforward calculations, it is easy to get

$$\Pi(\Omega, \mathbf{q}) = \frac{N\lambda^2}{16v_F v_\Delta}\frac{\Omega^2 + v_F^2 q_x^2}{q_1} + (q_x \leftrightarrow q_y), \tag{7}$$

where $q_1 = \sqrt{\Omega^2 + v_F^2 q_x^2 + v_\Delta^2 q_y^2}$. The polarization is linear in $|q|$, and dominates over the kinetic term q^2 at low energies. We can drop the q^2 term and write the effective nematic propagator as

$$D^{-1}(\Omega, \mathbf{q}) = r + \Pi(\Omega, \mathbf{q}). \tag{8}$$

The free fermion propagator will receive corrections due to fluctuation of nematic order parameter. After including these corrections, the complete propagator of Ψ_{1a} takes the following general form

$$G_{\Psi_{1a}}(\omega, \mathbf{k}) = \frac{1}{-i\omega A_0 + v_F k_x A_1\tau^z + v_\Delta k_y A_2\tau^x + m\tau^y}, \tag{9}$$

where $A_{0,1,2}$ are wave-function renormalizations and $m \equiv m(\omega, k_x, k_y)$ is a fermion gap. The mass gap term can never be generated as long as the fermion self-energy is calculated perturbatively by using the free fermion propagator $G^0(\omega, \mathbf{k})$. To examine the possibility of dynamical gap generation, we should instead utilize the following self-consistent DS equation

$$G_{\Psi_{1a}}^{-1}(\varepsilon, \mathbf{p}) = \left[G_{\Psi_{1a}}^0(\varepsilon, \mathbf{p})\right]^{-1} - \Sigma_{1a}(\varepsilon, \mathbf{p}) \tag{10}$$

where the self-energy is computed using the complete fermion propagator,

$$\Sigma_{1a}(\varepsilon, \mathbf{p}) = \lambda^2\int\frac{d\omega}{2\pi}\frac{d^2\mathbf{k}}{(2\pi)^2}\tau^x G_{\Psi_{1a}}(\omega, \mathbf{k})\tau^x \times\frac{1}{r + \Pi(\varepsilon - \omega, \mathbf{p} - \mathbf{k})}. \tag{11}$$

Substituting the complete propagator Eq.(9) into the DS equation, one can derive four self-consistently coupled equations of $A_{0,1,2}$ and m. Compared with the equation of m, the equations of $A_{0,1,2}$ contain an extra pre-factor of $1/N$, and are therefore relatively unimportant at large N. To the leading order of $1/N$ expansion, we assume that $A_0 = A_1 = A_2 = 1$, and then obtain a gap equation,

$$m(\varepsilon, p_x, p_y) = \lambda^2\int\frac{d\omega}{2\pi}\frac{d^2\mathbf{k}}{(2\pi)^2}\frac{m(\omega, k_x, k_y)}{\omega^2 + v_F^2 k_x^2 + v_\Delta^2 k_y^2 + m^2(\omega, k_x, k_y)} \times\frac{1}{r + \Pi(\epsilon - \omega, \mathbf{p} - \mathbf{k})}. \tag{12}$$

If this equation has only vanishing solution, $m = 0$, then the nematic fluctuation can not open any gap. A fermion gap is dynamically generated once this equation develops a nontrivial solution, $m \neq 0$. Due to the anisotropic nature of nematic fluctuation, the integrations over ω, k_x, and k_y have to be performed separately, which greatly increases the difficulty of numerical computations. For simplicity, here we consider the isotropic limit, $v_F = v_\Delta$. In this case, the dynamical gap becomes $m(\varepsilon, |\mathbf{p}|)$, and the polarization is simplified to

$$\Pi(\Omega, \mathbf{q}) = \frac{N\lambda^2}{16v_F^2}\frac{2\Omega^2 + v_F^2|\mathbf{q}|^2}{\sqrt{\Omega^2 + v_F^2|\mathbf{q}|^2}}. \tag{13}$$

We first consider the nematic QCP and take $r = 0$, then the pre-factor λ^2 on the right-hand side of Eq.(12) cancels exactly the factor λ^2 appearing in polarization $\Pi(\epsilon - \omega, \mathbf{p} - \mathbf{k})$. As a result, the gap equation is actually independent on coupling constant λ, and the critical



FIG. 1: (a) Relationship between dynamical gap $m(0,0)$ and fermion flavor N at nematic QCP, $r=0$; (b) Dependence of $m(0,0)$ on tuning parameter r for $N=2$. Dynamical gap is destroyed once r exceeds certain critical value r_c.

point of dynamical gap generation is solely determined by flavor N. After numerical computations, we find a critical fermion flavor $N_c \approx 2.4$ at $r=0$. The nodal qps remain gapless, $m=0$, when $N > N_c$, but acquire a nonzero dynamical gap m when $N < N_c$. Fig.1(a) presents the dependence of gap $m(\omega=0,\mathbf{k}=0)$ on flavor N. It is clear that the dynamical gap decreases very rapidly as flavor N grows, and vanishes continuously as $N \to N_c$. It is also interesting to examine how the tuning parameter r affects the gap generation. Actually, if we move away from the nematic QCP, r becomes finite and the nematic fluctuation is no longer critical. For $r \neq 0$, the coupling parameter λ cannot be exactly canceled, but one can absorb it into r by taking $r \to r/\lambda^2$. We find that the dynamical gap is significantly suppressed by growing r, and completely destroyed once r exceeds certain critical value r_c, which is shown in Fig.1(b). We therefore conclude that the dynamical gap generation is mediated by the critical fluctuation of nematic order parameter, and exists only in the vicinity of the nematic QCP.

Now let us discuss the physical meaning of the dynamical gap. The nonzero gap m of nodal qps generates an extra term which should be added to the Hamiltonian,

$$\begin{aligned} H_{\mathrm{m}} &= \int \frac{d^2\mathbf{k}}{(2\pi)^2}\{m(\Psi_{1a}^{\dagger}\tau^{y}\Psi_{1a} + \Psi_{2a}^{\dagger}\tau^{y}\Psi_{2a})\} \\ &= \int \frac{d^2\mathbf{k}}{(2\pi)^2}\left\{im\left[\left(f_{3\downarrow}^{\dagger}f_{1\uparrow}^{\dagger} + f_{3\downarrow}f_{1\uparrow}\right)\right.\right. \\ &\quad \left.\left. + \left(f_{1\downarrow}^{\dagger}f_{3\uparrow}^{\dagger} + f_{1\downarrow}f_{3\uparrow}\right) + (1 \leftrightarrow 2, 3 \leftrightarrow 4)\right\}\right. . \end{aligned}$$

One can see that such term corresponds to the formation of singlet Cooper pairs between the gapless qps excited from opposite nodes. We therefore have obtained a secondary, nematic-order driven, *is*-wave superconducting instability on top of the pure $d_{x^2-y^2}$ superconductivity.



FIG. 2: Schematic phase diagram. x_c denotes the nematic QCP. The shadowed region around x_c represents the emergent fully-gapped $d_{x^2-y^2}+is$ superconducting dome within a larger, pure $d_{x^2-y^2}$ superconducting dome.

As already pointed out, this dynamical gap is opened only when r is zero or very small, and is rapidly destroyed as r increases, which implies that the secondary *is*-wave superconductivity is achieved only in the close vicinity of nematic QCP. Upon approaching the nematic QCP, there is a zero-temperature phase transition from a pure $d_{x^2-y^2}$ superconducting state to a fully gapped $d_{x^2-y^2}+is$ superconducting state. At finite temperature, $T \neq 0$, the critical nematic fluctuation is weakened strongly due to the thermal screening effects, hence the dynamical nodal gap cannot survive at high temperatures. According to these analysis, we can infer that a small $d_{x^2-y^2}+is$ superconducting dome emerges around the nematic QCP, which is schematically shown in Fig. 2. It is interesting to notice that such nematic fluctuation-driven superconducting dome is analogous to that is formed on the border of an antiferromagnetic quantum critical point in some heavy fermion compounds [21].

Now we discuss the consequences of nonzero dynamical gap. For simplicity, let us assume a constant gap m, which gives rise to a new polarization function

$$\begin{aligned} \Pi(\mathbf{q},\Omega) &= \frac{N\lambda^2}{2\pi v_F v_\Delta}\frac{\Omega^2 + v_F^2 q_x^2}{q_1}\left[\frac{1}{2}\frac{m}{q_1} + \left(\frac{1}{4} - \frac{m^2}{q_1^2}\right)\right. \\ &\quad \left. \times \arcsin\left(\frac{q_1}{\sqrt{4m^2+q_1^2}}\right)\right] + (q_x \leftrightarrow q_y). \end{aligned}$$

In the low energy limit, it takes the simplified form $\Pi(\mathbf{q},\Omega) \approx \frac{N\lambda^2}{12\pi v_F v_\Delta m}(2\Omega^2 + v_F^2|\mathbf{q}|^2)$. Using this polarization, it is easy to obtain the following effective potential [22] between two massive nodal qps,

$$V(\mathbf{R}) \propto \int \frac{d^2\mathbf{q}}{(2\pi)^2}\frac{e^{i\mathbf{q}\cdot\mathbf{R}}}{\Pi(\mathbf{q})} \propto \frac{6v_\Delta}{N\lambda^2 v_F} m \ln(mR). \qquad (14)$$

This potential grows logarithmically as the distance R between two fermions is increasing, so the gapped fermions are weakly confined. Such a gap-induced confinement of fermions is similar to that in QED$_3$ [22].



FIG. 3: Momentum dependence of fermion velocities. Blue solid line: $\delta = 1$ and $m = 0$; Red Solid line: $\delta = 0.1$ and $m = 0$; Blue dashed line: $\delta = 1$ and $m/v_F\Lambda = 10^{-6}$; Red dashed line: $\delta = 0.1$ and $m/v_F\Lambda = 10^{-6}$.

When the nodal qps are massless, their velocities are renormalized by the nematic fluctuation and thus become strongly momentum-dependent, $v_{F,\Delta} \rightarrow v_{F,\Delta}(k)$. Both of $v_{F,\Delta}(k)$ vanish as $k \rightarrow 0$, but $v_\Delta(k)$ vanishes much faster than $v_F(k)$ [23], leading to extreme anisotropy, i.e., $\delta(k) = v_\Delta(k)/v_F(k) \rightarrow 0$ [15–19]. Nevertheless, once a finite gap m is opened, there is no longer strong renormalization of velocities. As shown in Fig. 3, the velocities $v_{F,\Delta}$ are still k-dependent and decrease with decreasing k at high energies, but saturate to finite values once k is smaller than the scale set by m. Therefore, the singular velocity renormalization and the extreme anisotropy are both prevented by the dynamical nodal gap m.

The dynamical gap also has important influence on observable quantities. For example, the density of state of nodal qps is linear in energy, $\rho(\omega) \propto |\omega|$ when $m = 0$, but becomes $\rho(\omega) \propto |\omega|\theta(|\omega| - m)$ when $m \neq 0$, which vanishes for $|\omega| < m$. Accordingly, the specific heat is strongly suppressed as $C_V \propto m^4/T^2 \exp(-m/T)$ in the temperature $T \ll m$. In addition, for finite m, the low temperature thermal conductivity is of the form [24], $\frac{\kappa}{T} = \frac{k_B^2}{3}(\frac{v_F}{v_\Delta} + \frac{v_\Delta}{v_F})\frac{\Gamma^2}{\Gamma^2+m^2}$, where Γ is impurity scattering rate. If $m = 0$, thermal conductivity is universal and impurity independent, i.e., $\frac{\kappa}{T} = \frac{k_B^2}{3}(\frac{v_F}{v_\Delta} + \frac{v_\Delta}{v_F})$ [25]. In contrast, for finite m, thermal conductivity is suppressed.

The nematic state is indeed equivalent to a superconducting state with $d_{x^2-y^2} + s$ gap, which was pointed out in Ref. [13]. Therefore, in the vicinity of a quantum critical point between a pure $d_{x^2-y^2} + s$ superconducting state and a $d_{x^2-y^2} + s$ superconducting state, the singular fluctuation of s-wave order parameter can also leads to a fully gapped $d_{x^2-y^2} + is$ superconducting state, provided that fermion flavor N is sufficiently small.

According to our results, it turns out that the fermion flavor N is a crucial parameter which determines the low-energy behaviors caused by the nematic order. When $N > N_c$, the non-perturbative effects of nematic fluctuation is unimportant, so one can trust the results obtained by perturbative calculations, such as extreme anisotropy [15] and other unusual properties [14, 16–20]. If $N < N_c$, however, the non-perturbative effects become significant, and can drive additional is-wave superconducting pairing between the originally gapless nodal qps. Our leading-order calculations found that $N_c \approx 2.4$, which is larger than physical flavor $N = 2$. It should be interesting to study how N_c is quantitatively affected by high order corrections in the future. In any case, our prediction of a nematic order-induced $d_{x^2-y^2} + is$ superconducting dome is novel and would shed light on the investigation of nematic order in correlated electron systems.

G.Z.L. acknowledges support by the National Natural Science Foundation of China under grant No. 11074234.

[1] S. A. Kivelson *et. al.*, Nature (London), **393**, 550 (1998).
[2] S. A. Kivelson *et. al.*, Rev. Mod. Phys. **75**, 1201 (2003).
[3] E. Fradkin *et. al.*, Annu. Rev. Condens. Matter Phys. **1**, 153 (2010); E. Fradkin, arXiv:1004.1104.
[4] M. Vojta, Adv. Phys. **58**, 699 (2009).
[5] Y. Ando *et. al.*, Phys. Rev. Lett. **88**, 137005 (2002).
[6] V. Hinkov *et. al.*, Science **319**, 597 (2008).
[7] R. Daou *et. al.*, Nature(London) **463**, 519 (2010).
[8] M. J. Lawler *et. al.*, Nature(London) **466**, 347 (2010).
[9] T.-M. Chuang *et. al.*, Science **327**, 181 (2010).
[10] R. Okazaki *et. al.*, Science **331**, 439 (2011).
[11] R. A. Borzi *et. al.*, Science **315**, 214(2007).
[12] K. B. Cooper *et. al.*, Phys. Rev. B **65**, 241313 (2002).
[13] M. Vojta, Y. Zhang, and S. Sachdev, Phys. Rev. B **62**, 6721 (2000); Int. J. Mod. Phys. B **14**, 3719 (2000).
[14] E.-A. Kim *et. al.*, Phys. Rev. B **77**, 184514 (2008).
[15] Y. Huh and S. Sachdev, Phys. Rev. B **78**, 064512 (2008).
[16] C. Xu, Y. Qi, and S. Sachdev, Phys. Rev. B **78**, 134507 (2008).
[17] L. Fritz and S. Sachdev, Phys. Rev. B **80**, 144503 (2009).
[18] J. Wang, G.-Z. Liu, and H. Kleinert, Phys. Rev. B **83**, 214503 (2011).
[19] G.-Z. Liu, J.-R. Wang, and J. Wang, Phys. Rev. B **85**, 174525 (2012).
[20] J. Wang and G.-Z. Liu, arXiv:1205.6164.
[21] H. v. Löhneysen *et al.*, Rev. Mod. Phys. **79**, 1015 (2007); O. Stockert *et al.*, J. Phys. Soc. Jpn. **81**, 011001 (2012).
[22] P. Maris, Phys. Rev. D **52**, 6087(1995).
[23] Expressions of $v_{F,\Delta}^{\mathrm{R}}(\omega, \mathbf{k})$ is too complicated to be shown here, but can be found in Refs. [15, 18].
[24] V. P. Gusynin and V. A. Miransky, Eur. Phys. J. **37**, 363 (2004).
[25] A. C. Durst and P. A. Lee, Phys. Rev. B **62**, 1270 (2000).